



**SCHOOL OF CHEMISTRY
UNIVERSITY OF HYDERABAD**



SEMINAR

Name of the Speaker	Prof. Tricia Naicker, University of KwaZulu-Natal (UKZN), South Africa
Title of the Talk	In Vitro and In Vivo Development of β - Lactam-Metallo- β -Lactamase Inhibitors
Date & Time	23.09.2026 (Wednesday) 4 PM
Venue	Seminar Hall, SoC

ALL ARE INVITED

Date: 15/09/2026

Faculty In-charge

Abstract:

In Vitro and In Vivo Development of β -Lactam-Metallo- β -Lactamase Inhibitors

β -lactams are the most prescribed class of antibiotics due to their potent, broad-spectrum antimicrobial activities. However, alarming rates of antimicrobial resistance now threaten the clinical relevance of these drugs, especially for the carbapenem-resistant Enterobacterales expressing metallo- β -lactamases (MBLs). Antimicrobial agents that specifically target these enzymes to restore the efficacy of last resort β -lactam drugs, that is, carbapenems, are therefore desperately needed. Herein, we present a cyclic zinc chelator covalently attached to a β -lactam scaffold (cephalosporin), that is, BP1. Observations from in vitro assays (with seven MBL expressing bacteria from different geographies) have indicated that BP1 restored the efficacy of meropenem to ≤ 0.5 mg/L, with sterilizing activity occurring from 8 h postinoculation. Furthermore, BP1 was nontoxic against human hepatocarcinoma cells (IC₅₀ $\approx$ 1000 mg/L) and exhibited a potency of (Ki app) 24.8 and 97.4 μ M against Verona integron-encoded MBL (VIM-2) and New Delhi metallo β -lactamase (NDM-1), respectively. There was no inhibition observed from BP1 with the human zinc-containing enzyme glyoxylase II up to 500 μ M. Preliminary molecular docking of BP1 with NDM-1 and VIM-2 sheds light on BP1's mode of action. In *Klebsiella pneumoniae* NDM infected mice, BP1 coadministered with meropenem was efficacious in reducing the bacterial load by ≈ 3 log₁₀ units' postinfection. The findings herein propose a favorable therapeutic combination strategy that restores the activity of the carbapenem antibiotic class and complements the few MBL inhibitors under development, with the ultimate goal of curbing antimicrobial resistance.